N.S.W.

Collection no. 99 — Moruya Dist.,

K.H.L. Key — 19–26.XII.1960.

19/12. 68 443.4 Potato Point.

68445.2 Maculata, paniculata & a stringybark to 50ft. with Acacia sp. to 15ft., Macrozamia, Casuarina ? tomulosa, & where thinned Cassinia sp. & a few other sclerophyllous shrubs. Acrida (j.), Gastr. (j.).
Eucalyptus australasica, Rhipidura moderata (j.), Cirphula pyrrhocnemis, Cryptoboa ruficollis chrysophonus, Bednura musgraveii, Tettix sp.

449.7. Bodalla. Proceed north.

453.2. Scabra, gummifera, pilularis, & occ. longifolia, all to 60ft, with a phyllodineous wattle to 15ft, normal wattle & Albizzia occ. to 10ft, occ. Cassinia & Macrozamia to 4ft. Themeda australis & Imperata to 18in, occ. microphyllous shrubs.
Morgoa amicula

Two other stops were made, in moderately favourable habitats for amicula, between Potato Point

+ Bodalla, but no amiculi was found. The site 453.2 is the first favourable one north of Bodalla, + is less than a mile from the southernmost previous locality for amiculi, forming part of the same system of ridges.

21/12.

68534. Moruya Heads. Took road to Moruya.

536. Maculata, scabra, gummifera, Phaulacridium vittatum, Coryphistes nigricosta, paniculata, longifolia, with Cassinia Paraletta argillacea, Petrin sp., Psednura microgravii to 6ft, Daviesia sp. + other small Leionema australasiae. shrubs + Macrozamia, all to 4ft, scattered Themeda australis + forbs. Gastr. (j.), Oed., Acrida (j.). Partly cleared + burned.

No habitats likely to harbour amiculi were seen anywhere along the Congo Park, Meringo Point, or Bingie roads.

Seen at Broulee, Dec. 1960: Phaulacridium vittatum, Gastr., Oed., Acrida (j.).